

FORMULAS GIVING PRIME NUMBERS UNDER CRAMÉR'S CONJECTURE

BAKIR FARHI

Département de Mathématiques, Université du Maine,
Avenue Olivier Messiaen, 72085 Le Mans Cedex 9, France.
Bakir.Farhi@univ-lemans.fr

ABSTRACT. Under Cramer's conjecture concerning the prime numbers, we prove that for any $\xi > 1$, there exists a real $A = A(\xi) > 1$ for which the formula $[A^{n^\xi}]$ (where $[.]$ denotes the integer part) gives a prime number for any positive integer n . Under the same conjecture, we also prove that for any $\varepsilon > 0$, there exists a real $B = B(\varepsilon) > 0$ for which the formula $[B.n!^{2+\varepsilon}]$ gives a prime number for any sufficiently large positive integer n .

MSC: 11A41.

Keywords: Prime numbers; Cramér's conjecture.

1. INTRODUCTION

Throughout this article, we will let $[x]$ denote the integer part of a giving real number x ; also, we will let $(p_n)_{n \in \mathbb{N}}$ denote the sequence of all prime numbers and we will put $\Delta p_n := p_{n+1} - p_n$ for any $n \in \mathbb{N}$. Further, if A is a subset of $\mathbb{R}$ and x is a real number, we will let $A+x$ denote the subset of $\mathbb{R}$ defined by: $A+x := \{a+x | a \in A\}$.

In [4], W. H. Mills proved the existence of an absolute constant $A > 1$ for which $[A^{3^n}]$ is a prime number for any positive integer n and in [6], E. M. Wright proved the existence of an absolute constant $\alpha > 0$ for which the infinite sequence $[\alpha], [2^\alpha], [2^{2^\alpha}], \dots$ is composed of prime numbers. Let us describe the method used by these two authors. They start from an upper bound of Δp_n as a function of p_n . Such upper bound allows to construct an increasing function h (more or less elementary, according to the used upper bound of Δp_n) such that between any two consecutive terms of the sequence $(h(n))_n$, there is at least one prime number. Setting $f_n := h \circ \dots \circ h$ (where h is applied n times), they deduce from the last fact, the existence of a real constant A for which the sequence $([f_n(A)])_n$ is consisted of prime numbers.

By this method, Wright uses the upper bound $\Delta p_n \leq p_n$ which is nothing else than Bertrand's postulate and Mills uses Ingham's upper bound: $\Delta p_n \leq p_n^{\frac{5}{8}+\varepsilon}$

(which holds for any sufficiently large n compared to a given $\varepsilon > 0$). The functions h which derive from these upper bounds are $h(x) = 2^x$ for Wright and $h(x) = x^3$ for Mills. Then the Theorems of [4] and [6] follow.

Notice that the more the upper bound of Δp_n is refined, the more the function h is small and the more the obtained sequence of prime numbers grows slowly (remark for instance that the sequence of Mills grows slower than the Wright's one). From this fact, in order to have a sequence of prime numbers which grows more slow again, we must use upper bounds more refined for Δp_n . But up to now even the powerful Riemann hypothesis gives only the estimate $\Delta p_n = O(p_n^{1/2} \log p_n)$. A famous conjecture (which is little too strong than this last estimate) states that between two consecutive squares, there is always a prime number (see [2]). So, according to this conjecture, the function $h(x) = x^2$ is admissible in the method described above, which we permits to conclude the existence of a constant $B > 1$ for which $[B^{2^n}]$ is a prime number for any positive integer n . We thus obtain (via this conjecture), a sequence of prime numbers growing slower than the Mills'one.

By leaning on heuristic and probabilistic arguments, H. Cramér [1] was leaded to conjecture that we have $\Delta p_n = O(\log^2 p_n)$; further it's known that $\Delta p_n = O(\log p_n)$ cannot hold (see [5]). Thus, by taking in the method described above $h(x) = c \log^2 x$ ($c > 0$), we obtain (via Cramér's conjecture) sequences of prime numbers having explicit form and growing much slower than the Mills'one. The inconvenient in this application is that the explicit form in question $[f_n(A)]$ is not elementary, because f_n doesn't have (in this case) a simple expression as function of n .

To cope with this problem, we were leaded to generalize Mills' method by considering instead of one function h , a sequence of functions $(h_m)_m$ and in this situation f_n is rather the composition of n functions $h_0, \dots, h_{n-1}$. This allows fundamentally to give for f_n the form which we want, then if we set $h_n := f_{n+1} \circ f_n^{-1}$, we have only to check whether it's true that for any n and any x sufficiently large (relative to n), the interval $[h_n(x), h_n(x+1) - 1[$ contains at least one prime number or not. In the affirmative case, we will deduce the existence of a real A for which the formula $[f_n(A)]$ gives a prime number for any positive integer n (see Theorem 1 and its proof).

Under a conjecture less strong than the Cramér's one, we derive from this generalization two new types of explicit formulas giving prime numbers. We also give other applications of our main result (outside the subject of prime numbers) and we conclude this article by some open questions related to the results which we obtain.

2. RESULTS

The main result of this article is the following

Theorem 1. Let $I =]a, b[$ (with $a, b \in \overline{\mathbb{R}}$, $a < b$) be an open interval of $\mathbb{R}$, n_0 be a non-negative integer and $(f_n)_{n \geq n_0}$ be a sequence of real functions which are differentiable and increasing on I .

We assume that the functions $\frac{f'_{n+1}}{f'_n}$ ($n \geq n_0$) are nondecreasing on I and that for all $x \in I$, the numerical sequence $(f_n(x))_{n \geq n_0}$ is increasing.

We also assume that there exists a real function g , nondecreasing on $\mathbb{R}$ and verifying:

$$(1) \quad g \circ f_{n+1}(x) \leq \frac{f'_{n+1}}{f'_n}(x) \quad (\forall n \geq n_0 \text{ and } \forall x \in I).$$

Then, for any sequence of integers $(u_n)_{n \in \mathbb{N}}$, verifying: $\limsup_{n \rightarrow +\infty} u_n = +\infty$,

$$(2) \quad u_{n+1} - u_n \leq g(u_n) - 1 \quad (\text{for all } n \text{ sufficiently large})$$

and whose one at least of the terms u_n (n satisfying (2)) belongs to $f_{n_0}(I) \cap (f_{n_0}(I) - 1)$, there exists a real $A \in I$ for which the sequence $([f_n(A)])_{n \geq n_0}$ is an increasing subsequence of $(u_n)_n$.

Proof: By shifting if necessary the sequence of functions $(f_n)_{n \geq n_0}$, we may assume that $n_0 = 0$ and by shifting if necessary the sequence $(u_n)_n$, we may assume that we have (more generally than (2)):

$$(2') \quad u_{n+1} - u_n \leq g(u_n) - 1 \quad (\text{for all } n \in \mathbb{N}).$$

We begin this proof by some remarks and preliminary notations which allow to lighten better the situation of the Theorem.

Giving $n \in \mathbb{N}$, since the function f_n is assumed differentiable (so continuous) and increasing on $I =]a, b[$, then it's a bijection from I into $f_n(I) =]\lim_{x \rightarrow a} f_n(x), \lim_{x \rightarrow b} f_n(x)[=]\lambda_n, \mu_n[$, where $\lambda_n := \lim_{x \rightarrow a} f_n(x)$ and $\mu_n := \lim_{x \rightarrow b} f_n(x)$ (λ_n and μ_n belong to $\overline{\mathbb{R}}$).

Now, let us introduce the following functions:

$$h_n :]\lambda_n, \mu_n[\longrightarrow]\lambda_{n+1}, \mu_{n+1}[\quad (\forall n \in \mathbb{N}).$$

$$h_n := f_{n+1} \circ f_n^{-1}$$

Giving $n \in \mathbb{N}$, since (from the hypothesis of the Theorem), the functions f_n and f_{n+1} are differentiable and increasing on I , then the function h_n is differentiable and increasing on $] \lambda_n, \mu_n[$. Further, the hypothesis of the Theorem concerning the increase of the numerical sequences $(f_n(x))_n$ ($x \in I$) amounts to:

$$(3) \quad h_n(x) > x \quad (\forall n \in \mathbb{N} \text{ and } \forall x \in]\lambda_n, \mu_n[).$$

Next, let us show that for any $n \in \mathbb{N}$, the function h_n is convex on $] \lambda_n, \mu_n[$. To do this, we check that the derivative h'_n of each function h_n ($n \in \mathbb{N}$) is nondecreasing on the interval $] \lambda_n, \mu_n[$. Giving $n \in \mathbb{N}$, we have:

$$h'_n = (f_{n+1} \circ f_n^{-1})' = (f_n^{-1})' \cdot f'_{n+1} \circ f_n^{-1} = \frac{f'_{n+1} \circ f_n^{-1}}{f'_n \circ f_n^{-1}} = \left(\frac{f'_{n+1}}{f'_n} \right) \circ f_n^{-1}.$$

Since (from the hypothesis of the Theorem), the function $\frac{f'_{n+1}}{f'_n}$ is nondecreasing on I and the function f_n^{-1} is increasing on $f_n(I) =]\lambda_n, \mu_n[$ (because f_n is increasing on I), then (as a composite of two nondecreasing functions), the function h'_n is nondecreasing on $]\lambda_n, \mu_n[$. So the function h_n is effectively convex on $]\lambda_n, \mu_n[$.

The rest of this proof consists of the three following steps:

1st Step: We are going to show that we have:

$$(4) \quad g \circ h_n(y) \leq h_n(y+1) - h_n(y) \quad (\forall n \in \mathbb{N} \text{ and } \forall y \in]\lambda_n, \mu_n - 1[).$$

(We will see further that the interval $]\lambda_n, \mu_n - 1[$ is never empty).

Let $n \in \mathbb{N}$ and $y \in]\lambda_n, \mu_n - 1[$ fixed and set $x := f_n^{-1}(y)$.

The convexity of h_n on $]\lambda_n, \mu_n[$, proved above, implies that we have:

$$h_n(u) \geq h'_n(t)(u-t) + h_n(t) \quad (\text{for all } t, u \in]\lambda_n, \mu_n[).$$

By taking in this last inequality $t = y$ and $u = y+1$, we obtain:

$$\begin{aligned} h_n(y+1) - h_n(y) &\geq h'_n(y) \\ &= \left(\frac{f'_{n+1}}{f'_n} \right)(x) \quad (\text{because } h'_n = \frac{f'_{n+1}}{f'_n} \circ f_n^{-1} \text{ and } x = f_n^{-1}(y)) \\ &\geq g \circ f_{n+1}(x) \quad (\text{from the hypothesis (1) of the Theorem}) \\ &= g \circ f_{n+1} \circ f_n^{-1}(y) \\ &= g \circ h_n(y). \end{aligned}$$

The relation (4) follows.

2nd Step: We are going to construct an increasing sequence $(k_n)_{n \in \mathbb{N}}$ of non-negative integers such that the subsequence of $(u_n)_n$ with general term $v_n = u_{k_n}$ satisfies:

$$\begin{cases} v_n \in]\lambda_n, \mu_n - 1[\\ \text{and} \\ h_n(v_n) \leq v_{n+1} < h_n(v_n + 1) - 1 \end{cases} \quad (\forall n \in \mathbb{N}).$$

We proceed by induction as follows:

- We pick $k_0 \in \mathbb{N}$ such that $u_{k_0} \in f_0(I) \cap (f_0(I) - 1) =]\lambda_0, \mu_0 - 1[$. Notice that the existence of such integer k_0 is an hypothesis of the Theorem.
- If, for some $n \in \mathbb{N}$, an integer $k_n \in \mathbb{N}$ is chosen such that $u_{k_n} \in]\lambda_n, \mu_n - 1[$, let:

$$X_n := \{k \in \mathbb{N} \mid k > k_n \text{ and } u_k \geq h_n(u_{k_n})\}.$$

From the hypothesis $\limsup_{n \rightarrow +\infty} u_n = +\infty$, this subset X_n of $\mathbb{N}$ is nonempty, it thus admits a smallest element which we take as the choice of k_{n+1} . So, we have:

$$k_{n+1} > k_n, \quad u_{k_{n+1}} \geq h_n(u_{k_n}) \text{ and } k_{n+1} - 1 \notin X_n.$$

We claim that the facts “ $k_{n+1} > k_n$ ” and “ $k_{n+1} - 1 \notin X_n$ ” imply:

$$(5) \quad u_{k_{n+1}-1} < h_n(u_{k_n}).$$

Indeed: either $k_{n+1} = k_n + 1$ in which case we have $u_{k_{n+1}-1} = u_{k_n} < h_n(u_{k_n})$ (from (3)); or $k_{n+1} > k_n + 1$, that is $k_{n+1} - 1 > k_n$, but since $k_{n+1} - 1 \notin X_n$, we are forced to have (also in this case) $u_{k_{n+1}-1} < h_n(u_{k_n})$ as required.

It follows that we have:

$$\begin{aligned} u_{k_{n+1}} &\leq u_{k_{n+1}-1} + g(u_{k_{n+1}-1}) - 1 && \text{(from (2'))} \\ &< h_n(u_{k_n}) + g \circ h_n(u_{k_n}) - 1 && \text{(by using (5) and the non-decrease of } g) \\ &\leq h_n(u_{k_n} + 1) - 1 && \text{(from (4)).} \end{aligned}$$

Hence:

$$u_{k_{n+1}} < h_n(u_{k_n} + 1) - 1.$$

We thus have:

$$h_n(u_{k_n}) \leq u_{k_{n+1}} < h_n(u_{k_n} + 1) - 1.$$

Finally, since the function h_n takes its values in $] \lambda_{n+1}, \mu_{n+1} [$, then the last double inequality does show that $u_{k_{n+1}} \in] \lambda_{n+1}, \mu_{n+1} - 1 [$. This ensures the good working of this induction process which gives the required sequence $(k_n)_n$.

Notice also that the subsequence $(v_n)_n$ of $(u_n)_n$ which we have just constructed is increasing, because for any $n \in \mathbb{N}$, we have: $v_{n+1} \geq h_n(v_n) > v_n$ (from (3)).

3rd step (conclusion): To conclude this proof, we will show the existence of a real $A \in I$ for which we have for any $n \in \mathbb{N}$: $v_n = [f_n(A)]$.

To do this, we introduce two real sequences $(x_n)_n$ and $(y_n)_n$, with terms in I , which we define by:

$$x_n := f_n^{-1}(v_n) \text{ and } y_n := f_n^{-1}(v_n + 1) \quad (\forall n \in \mathbb{N}).$$

Since the functions f_n are increasing, we have $x_n < y_n$ for all $n \in \mathbb{N}$. We claim that the sequence $(x_n)_n$ is nondecreasing and that the sequence $(y_n)_n$ is decreasing. Indeed, for any $n \in \mathbb{N}$, we have:

$$x_n = f_n^{-1}(v_n) = f_{n+1}^{-1} \circ h_n(v_n) \leq f_{n+1}^{-1}(v_{n+1}) = x_{n+1}$$

and

$$y_n = f_n^{-1}(v_n + 1) = f_{n+1}^{-1} \circ h_n(v_n + 1) > f_{n+1}^{-1}(v_{n+1} + 1) = y_{n+1}.$$

(Where in these last relations, we have just use the facts that f_{n+1}^{-1} is increasing and $h_n(v_n) \leq v_{n+1} < h_n(v_n + 1) - 1$).

The $[x_n, y_n]$ ($n \in \mathbb{N}$) are thus nested inclosed intervals of $\mathbb{R}$. Consequently their intersection is nonempty (according to the Cantor's intersection theorem). Pick A an arbitrary real belonging to this intersection, that is: $x_n \leq A \leq y_n$ for all $n \in \mathbb{N}$ (in particular $A \in I$). In fact A verifies even:

$$x_n \leq A < y_n \quad (\forall n \in \mathbb{N}),$$

because if $A = y_m$ for some $m \in \mathbb{N}$, we will have (from the decreasing of the sequence $(y_n)_n$): $A > y_{m+1}$, contradicting the inequality $A \leq y_{m+1}$.

It follows from the increase of the functions f_n that we have:

$$f_n(x_n) \leq f_n(A) < f_n(y_n) \quad (\forall n \in \mathbb{N}),$$

that is:

$$v_n \leq f_n(A) < v_n + 1 \quad (\forall n \in \mathbb{N}).$$

Then (since v_n is an integer for all $n \in \mathbb{N}$):

$$[f_n(A)] = v_n \quad (\forall n \in \mathbb{N}).$$

This completes the proof. ■

Remarks: Mills' theorem [4] can be find again by applying the above Theorem 1 for $I =]1, +\infty[$, $n_0 = 0$, $f_n(x) = x^{3^n}$ ($\forall n \in \mathbb{N}$ and $\forall x \in I$), $g(x) = x^{2/3}$ if $x > 0$ and $g(x) = 0$ if $x \leq 0$ and $(u_n)_n$ the sequence of the prime numbers. In this application, we check the relation (1) of Theorem 1 by a simple calculus and we deduce the relation (2) of the same Theorem from the Ingham's estimate quoted in the introduction. The remaining hypothesis of Theorem 1 are immediately verified.

The Wright's theorem [6] can be also find again by applying Theorem 1 for $I =]0, +\infty[$, $n_0 = 0$, $(f_n)_{n \in \mathbb{N}}$ the sequence of functions which is defined on I by: $f_0 = \text{Id}_I$ and $f_{n+1} = 2^{f_n}$ ($\forall n \in \mathbb{N}$), $g(x) = (\log 2)x$ ($\forall x \in \mathbb{R}$) and $(u_n)_n$ the sequence of the prime numbers. In order to check the relation (1) of Theorem 1, remark that we have for any $n \in \mathbb{N}$: $\frac{f'_{n+1}}{f'_n} = (\log 2)f_{n+1}$. As for the relation (2), it's (in this application) a consequence of the prime numbers theorem; but it can be obtained by using Chebyshev's elementary arguments (see [3]). The remaining hypothesis of Theorem 1 are immediately verified.

N. B.— It's interesting to remark that in the two above applications of Theorem 1, the sequence of functions $(h_n)_n$ introduced in the proof of the latter ($h_n := f_{n+1} \circ f_n^{-1}$) is constant. Indeed, for the first application, we have $h_n(x) = x^3$ ($\forall n \in \mathbb{N}$) and for the second one, we have $h_n(x) = 2^x$ ($\forall n \in \mathbb{N}$). As explained in the introduction, the fact to be able to take $(h_n)_n$ not constant is the crucial point of our approach. In the following, we are going to give some applications of Theorem 1 in which this sequence $(h_n)_n$ is not constant. If we admit the following Conjecture (which is less strong than the Cramér's one [1]), we obtain two new types of explicit sequences of prime numbers, which grow much more slow than the ones of Mills and Wright.

Conjecture 2. *there exists an absolute constant $k > 1$ such that:*

$$\Delta p_n = O((\log p_n)^k).$$

Under this Conjecture, we obtain by applying Theorem 1, the two following Corollaries:

Corollary 3. *Under the above Conjecture 2, there exists for all real $\xi > 1$, a real $A = A(\xi) > 1$ for which the sequence $([A^{n^\xi}])_{n \geq 1}$ is an increasing sequence of prime numbers.*

Proof: Let $\xi > 1$ be fixed, $k > 1$ be an admissible constant in Conjecture 2 and $a > 1$ be a real such that:

$$(6) \quad (\log x)^{k+1} \leq x^{1/2} \quad (\forall x > a)$$

$$(7) \quad (n+1)^{k+1} \leq a^{\frac{1}{2}n^{\xi-1}} \quad (\forall n \geq 1).$$

(a exists because $\lim_{x \rightarrow +\infty} \frac{(\log x)^{k+1}}{x^{1/2}} = 0$ and $\lim_{n \rightarrow +\infty} (n+1)^{\frac{2(k+1)}{n^{\xi-1}}} = 1$).

We apply Theorem 1 for $I =]a, +\infty[$, $n_0 = 1$, $f_n(x) = x^{n^\xi}$ ($\forall n \geq 1, \forall x \in I$), $g(x) = (\log x)^{k+1}$ if $x > 1$ and $g(x) = 0$ if $x \leq 1$ and $(u_n)_n$ the sequence of the prime numbers. Let us check the hypothesis of Theorem 1:

The functions f_n are clearly increasing and differentiable on I . We have $f'_n(x) = n^\xi x^{n^\xi-1}$ ($\forall n \geq 1, \forall x \in I$), then: $\frac{f'_{n+1}}{f'_n}(x) = (\frac{n+1}{n})^\xi x^{(n+1)^\xi - n^\xi}$ ($\forall n \geq 1, \forall x \in I$). We thus see that the functions $\frac{f'_{n+1}}{f'_n}$ ($n \geq 1$) are nondecreasing on I . Further, if x is a fixed real in I , the numerical sequence $(f_n(x))_{n \geq 1}$ is clearly increasing. Now, we have for any integer $n \geq 1$ and for any real $x \in I$:

$$\begin{aligned} g \circ f_{n+1}(x) &= (n+1)^{\xi(k+1)} (\log x)^{k+1} \\ &\leq a^{\frac{1}{2}\xi n^{\xi-1}} x^{1/2} && \text{(from (6) and (7))} \\ &\leq x^{\xi n^{\xi-1}} && \text{(because } x > a \text{ and } \xi n^{\xi-1} > 1) \\ &\leq x^{(n+1)^\xi - n^\xi} && \text{(because } \xi n^{\xi-1} \leq (n+1)^\xi - n^\xi) \\ &\leq \frac{f'_{n+1}}{f'_n}(x). \end{aligned}$$

The relation (1) of Theorem 1 follows.

Next, the relation (2) of Theorem 1 (related to this application) follows immediately from the conjecture 2 (admitted in this context). Finally, $f_{n_0}(I) \cap (f_{n_0}(I) - 1) =]a, +\infty[$ does contains prime numbers as large as we want. The hypothesis of Theorem 1 are thus all satisfied, so we can apply this latter to the present situation. The corollary 3 follows from this application. $\blacksquare$

Corollary 4. *Assume that Conjecture 2 is true and let $k > 1$ be an admissible constant in this Conjecture. Then, for any positive real ε , there exists an integer $n_0 = n_0(\varepsilon, k) \geq 1$ and a real $B = B(\varepsilon, k) > 0$ such that the sequence $([B \cdot n!^{k+\varepsilon}])_{n \geq n_0}$ is an increasing sequence of prime numbers.*

Proof: let ε be a fixed positive real. From Conjecture 2 (admitted with the constant $k > 1$), there exists a positive real c_k for which we have:

$$(8) \quad p_{n+1} - p_n \leq c_k (\log p_n)^k \quad (\forall n \in \mathbb{N}).$$

We apply Theorem 1 for $I =]1, 2[$, $n_0 \geq 2$ an integer (depending on k and ε) which we pick large enough that we have:

$$(9) \quad c_k ((k + \varepsilon)(n + 1) \log(n + 1) + \log 2)^k + 1 \leq (n + 1)^{k+\varepsilon} \quad (\forall n \geq n_0),$$

$f_n(x) = n!^{k+\varepsilon} x$ ($\forall n \geq n_0, \forall x \in I$), $g(x) = c_k (\log x)^k + 1$ if $x > 1$ and $g(x) = 1$ if $x \leq 1$ and $(u_n)_n$ the sequence of the prime numbers. In this situation, we can easily check that the hypothesis of Theorem 1 are all satisfied. We just note that the relation (1) follows from (9), the relation (2) follows from (8) and the last hypothesis of Theorem 1 concerning the sequence $(u_n)_n = (p_n)_n$ is a consequence of Bertrand's postulate. Corollary 4 follows from this application. ■

Apart from the context of the prime numbers, we have the following

Corollary 5. *Let $(u_n)_{n \in \mathbb{N}}$ be a sequence of integers such that: $1 \leq \limsup_{n \rightarrow +\infty} (u_{n+1} - u_n) < +\infty$. Then, we have:*

- (1) *For any positive real λ , there exists a real $A > 1$ for which the sequence $([\lambda A^n])_{n \geq 1}$ is an increasing subsequence of $(u_n)_n$.*
- (2) *For any real $A > \limsup_{n \rightarrow +\infty} (u_{n+1} - u_n) + 1$, there exists a positive real λ for which the sequence $([\lambda A^n])_{n \geq 1}$ is an increasing subsequence of $(u_n)_n$.*

Some open problems related to the preceding study: We ask (under or without Cramér's Conjecture) the following questions:

- (1) Does there exists a real number $A > 1$ for which $[A^n]$ is a prime number for any positive integer n ? (This corresponds to the case $\xi = 1$ which is excluded from Corollary 3).
- (2) More generally than (1), does there exists a couple of real numbers (λ, A) , with $\lambda > 0$, $A > 1$, for which $[\lambda A^n]$ is a prime number for any positive integer n ? (This is related to Corollary 5).
- (3) Does there exists a real number $B > 1$ for which $[B.n!^2]$ is a prime number for any sufficiently large non-negative integer n ? (This is related to Corollary 4).

REFERENCES

- [1] H. CRAMÉR. On the order of magnitude of the difference between consecutive prime numbers. *Acta Arith.*, **2** (1936), p. 23-46.
- [2] R. K. GUY. Unsolved problems in Number Theory. 2 nd ed., *Springer-Verlag*, New York, (1994).
- [3] G. H. HARDY & E. M. WRIGHT. The theory of numbers. *Oxford Univ. Press*, London, 5th ed, (1979).

- [4] W. H. MILLS. A prime-representing function. *Bull. Amer. Math. Soc*, **53** (1947), p. 604.
- [5] E. WESTZYNTHIUS. Über die Verteilung der Zahlen die zu den n ersten Primzahlen teilerfremd sind. *Comm. Phys. Math. Helsingfors*, (5), **25** (1931), p. 1-37.
- [6] E. M. WRIGHT. A prime-representing function. *Amer. Math. Monthly*, **58** (1951), p. 616-618.